

## Math 1600 Lecture 28, Section 2, 12 Nov 2014

### Announcements:

Today we review 4.3 and discuss Appendix C. Next class we finish 4.3 and start 4.4. **Read** Section 4.4 for next class. Work through recommended [homework questions](#). Exercises and solutions for Appendix C are posted on that page.

Next class: **course evaluations at start.**

**Final exam:** Monday, December 8, 9am to noon. See the [course home page](#) for final exam **conflict** policy. You should **immediately** notify the registrar or your Dean's office (and your instructor) of any conflicts! (Deadline Nov 21.)

**Tutorials:** Quiz 7 covers 4.2, the parts of Appendix D that we covered, and the part of 4.3 we finished Monday. No complex eigenvalues/roots.

**Help Centers:** Monday-Friday 2:30-6:30 in MC 106.

**Office hour:** Wednesday, 11:30-noon, MC103B.

### Brief review of last lecture:

The **characteristic polynomial** of a square matrix  $A$  is  $\det(A - \lambda I)$ , which is a polynomial in  $\lambda$ . The roots/zeros of this polynomial are the eigenvalues of  $A$ .

A root  $a$  of a polynomial  $f$  implies that  $f(x) = (x - a)g(x)$ . Sometimes,  $a$  is also a root of  $g(x)$ . Then  $f(x) = (x - a)^2 h(x)$ . The largest  $k$  such that  $(x - a)^k$  is a factor of  $f$  is called the **multiplicity** of the root  $a$  in  $f$ .

In the case of an eigenvalue, we call its multiplicity in the characteristic polynomial the **algebraic multiplicity** of this eigenvalue.

For example, if  $\det(A - \lambda I) = -(\lambda - 1)^2(\lambda - 2)$ , then  $\lambda = 1$  is an eigenvalue with algebraic multiplicity 2, and  $\lambda = 2$  is an eigenvalue with algebraic multiplicity 1.

We also define the **geometric multiplicity** of an eigenvalue  $\lambda$  to be the dimension of the corresponding eigenspace.

**Theorem 4.15:** The eigenvalues of a triangular matrix are the entries on its main diagonal (repeated according to their algebraic multiplicity).

**Theorem 4.17:** Let  $A$  be an  $n \times n$  matrix. The following are equivalent:

- a.  $A$  is invertible.
- b.  $A\vec{x} = \vec{b}$  has a unique solution for every  $\vec{b} \in \mathbb{R}^n$ .
- c.  $A\vec{x} = \vec{0}$  has only the trivial (zero) solution.
- d. The reduced row echelon form of  $A$  is  $I_n$ .
- f.  $\text{rank}(A) = n$
- g.  $\text{nullity}(A) = 0$
- h. The columns of  $A$  are linearly independent.
- i. The columns of  $A$  span  $\mathbb{R}^n$ .
- j. The columns of  $A$  are a basis for  $\mathbb{R}^n$ .
- k. The rows of  $A$  are linearly independent.
- l. The rows of  $A$  span  $\mathbb{R}^n$ .
- m. The rows of  $A$  are a basis for  $\mathbb{R}^n$ .
- n.  $\det A \neq 0$
- o. 0 is not an eigenvalue of  $A$

## Eigenvalues of powers and inverses

**Theorem 4.18:** If  $\vec{x}$  is an eigenvector of  $A$  with eigenvalue  $\lambda$ , then  $\vec{x}$  is an eigenvector of  $A^k$  with eigenvalue  $\lambda^k$ . This holds for each integer  $k \geq 0$ , and also for  $k < 0$  if  $A$  is invertible.

In contrast to some other recent results, this one is very useful computationally:

**Example 4.21:** Compute  $\begin{bmatrix} 0 & 1 \\ 2 & 1 \end{bmatrix}^{10} \begin{bmatrix} 5 \\ 1 \end{bmatrix}$ .

See last lecture for the method used, which is much faster than repeated matrix multiplication.

**Theorem:** If  $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m$  are eigenvectors of  $A$  corresponding to **distinct** eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_m$ , then  $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m$  are linearly independent.

## New material: Appendix C: Complex numbers

Sometimes a polynomial has complex numbers as its roots, so we need to learn a bit about them.

A **complex number** is a number of the form  $a + bi$ , where  $a$  and  $b$  are real numbers and  $i$  is a symbol such that  $i^2 = -1$ .

If  $z = a + bi$ , we call  $a$  the **real part** of  $z$ , written  $\operatorname{Re} z$ , and  $b$  the **imaginary part** of  $z$ , written  $\operatorname{Im} z$ .

Complex numbers  $a + bi$  and  $c + di$  are **equal** if  $a = c$  and  $b = d$ .

On board: sketch complex plane and various points.

**Addition:**  $(a + bi) + (c + di) = (a + c) + (b + d)i$ , like vector addition.

**Multiplication:**  $(a + bi)(c + di) = (ac - bd) + (ad + bc)i$ . (Explain.)

**Examples:**  $(1 + 2i) + (3 + 4i) = 4 + 6i$

$$\begin{aligned}(1 + 2i)(3 + 4i) &= 1(3 + 4i) + 2i(3 + 4i) = 3 + 4i + 6i + 8i^2 \\ &= (3 - 8) + 10i = -5 + 10i\end{aligned}$$

$$5(3 + 4i) = 15 + 20i$$

$$(-1)(c + di) = -c - di$$

The **conjugate** of  $z = a + bi$  is  $\bar{z} = a - bi$ . Reflection in real axis. We'll use this for division of complex numbers in a moment.

**Theorem (Properties of conjugates):** Let  $w$  and  $z$  be complex numbers. Then:

1.  $\overline{\bar{z}} = z$
2.  $\overline{w + z} = \bar{w} + \bar{z}$
3.  $\overline{wz} = \bar{w}\bar{z}$  (typo in text) (good exercise)
4. If  $z \neq 0$ , then  $\overline{w/z} = \bar{w}/\bar{z}$  (see below for division)
5.  $z$  is **real** if and only if  $\bar{z} = z$

The **absolute value** or **modulus**  $|z|$  of  $z = a + bi$  is

$$|z| = |a + bi| = \sqrt{a^2 + b^2}, \quad \text{the distance from the origin.}$$

**Examples:**  $|3| = 3$ ,  $|-3| = 3$ ,  $|\pm i| = 1$ ,  $|3 + 4i| = \sqrt{3^2 + 4^2} = 5$ .

Note that

$$z\bar{z} = (a + bi)(a - bi) = a^2 - abi + abi - b^2i^2 = a^2 + b^2 = |z|^2$$

This means that for  $z \neq 0$

$$\frac{z\bar{z}}{|z|^2} = 1 \quad \text{so} \quad z^{-1} = \frac{\bar{z}}{|z|^2}$$

This can be used to compute quotients of complex numbers:

$$\frac{w}{z} = \frac{w}{z} \frac{\bar{z}}{\bar{z}} = \frac{w\bar{z}}{|z|^2}.$$

**Example:**

$$\frac{-1 + 2i}{3 + 4i} = \frac{-1 + 2i}{3 + 4i} \frac{3 - 4i}{3 - 4i} = \frac{5 + 10i}{3^2 + 4^2} = \frac{5 + 10i}{25} = \frac{1}{5} + \frac{2}{5}i$$

**Theorem (Properties of absolute value):** Let  $w$  and  $z$  be complex numbers.

Then:

1.  $|z| = 0$  if and only if  $z = 0$ .
2.  $|\bar{z}| = |z|$
3.  $|wz| = |w||z|$  (good exercise!)
4. If  $z \neq 0$ , then  $|w/z| = |w|/|z|$ . In particular,  $|1/z| = 1/|z|$ .
5.  $|w + z| \leq |w| + |z|$ .

## Polar Form

A complex number  $z = a + bi$  can also be expressed in **polar coordinates**  $(r, \theta)$ , where  $r = |z| \geq 0$  and  $\theta$  is such that

$$a = r \cos \theta \quad \text{and} \quad b = r \sin \theta \quad (\text{sketch})$$

Then

$$z = r \cos \theta + (r \sin \theta)i = r(\cos \theta + i \sin \theta)$$

To compute  $\theta$ , note that

$$\tan \theta = \sin \theta / \cos \theta = b/a.$$

But this doesn't pin down  $\theta$ , since  $\tan(\theta + \pi) = \tan \theta$ . You must choose  $\theta$  based on what quadrant  $z$  is in. There is a unique correct  $\theta$  with  $-\pi < \theta \leq \pi$ , and this is

called the **principal argument** of  $z$  and is written  $\text{Arg } z$  (or  $\arg z$ ).

**Examples:** If  $z = 1 + i$ , then  $r = |z| = \sqrt{1^2 + 1^2} = \sqrt{2}$ . By inspection,  $\theta = \pi/4 = 45^\circ$ . We also know that  $\tan \theta = 1/1 = 1$ , which gives  $\theta = \pi/4 + k\pi$ , and  $k = 0$  gives the right quadrant.

We write  $\text{Arg } z = \pi/4$  and  $z = \sqrt{2}(\cos \pi/4 + i \sin \pi/4)$ .

If  $w = -1 - i$ , then  $r = \sqrt{2}$  and by inspection  $\theta = -3\pi/4 = -135^\circ$ . We *still* have  $\tan \theta = -1/-1 = 1$ , which gives  $\theta = \pi/4 + k\pi$ , but now we must take  $k$  odd to land in the right quadrant. Taking  $k = -1$  gives the principal argument:

$$\text{Arg } w = -3\pi/4 \quad \text{and} \quad w = \sqrt{2}(\cos(-3\pi/4) + i \sin(-3\pi/4)).$$

## Multiplication and division in polar form

Let

$$z_1 = r_1(\cos \theta_1 + i \sin \theta_1) \quad \text{and} \quad z_2 = r_2(\cos \theta_2 + i \sin \theta_2).$$

Then

$$\begin{aligned} z_1 z_2 &= r_1 r_2 (\cos \theta_1 + i \sin \theta_1)(\cos \theta_2 + i \sin \theta_2) \\ &= r_1 r_2 [(\cos \theta_1 \cos \theta_2 - \sin \theta_1 \sin \theta_2) + i(\sin \theta_1 \cos \theta_2 + \cos \theta_1 \sin \theta_2)] \\ &= r_1 r_2 [\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2)] \end{aligned}$$

So

$$|z_1 z_2| = |z_1| |z_2| \quad \text{and} \quad \text{Arg}(z_1 z_2) = \text{Arg } z_1 + \text{Arg } z_2$$

(up to multiples of  $2\pi$ ). Sketch on board. See also Example C.4.

In particular, if  $z = r(\cos \theta + i \sin \theta)$ , then  $z^2 = r^2(\cos(2\theta) + i \sin(2\theta))$ . It follows that the two **square roots** of  $z$  are

$$\pm \sqrt{r}(\cos(\theta/2) + i(\sin \theta/2))$$

## The remaining material is for your interest only:

Repeating this argument gives:

**Theorem (De Moivre's Theorem):** If  $z = r(\cos \theta + i \sin \theta)$  and  $n$  is a positive

integer, then

$$z^n = r^n (\cos(n\theta) + i \sin(n\theta))$$

When  $r \neq 0$ , this also holds for  $n$  negative. In particular,

$$\frac{1}{z} = \frac{1}{r} (\cos \theta - i \sin \theta).$$

**Example C.5:** Find  $(1 + i)^6$ .

**Solution:** We saw that  $1 + i = \sqrt{2}(\cos(\pi/4) + i \sin(\pi/4))$ . So

$$\begin{aligned} (1 + i)^6 &= (\sqrt{2})^6 (\cos(6\pi/4) + i \sin(6\pi/4)) \\ &= 8(\cos(3\pi/2) + i \sin(3\pi/2)) \\ &= 8(0 + i(-1)) = -8i \end{aligned}$$

## $n$ th roots

De Moivre's Theorem also lets us compute  $n$ th roots:

**Theorem:** Let  $z = r(\cos \theta + i \sin \theta)$  and let  $n$  be a positive integer. Then  $z$  has exactly  $n$  distinct  $n$ th roots, given by

$$r^{1/n} \left[ \cos \left( \frac{\theta + 2k\pi}{n} \right) + i \sin \left( \frac{\theta + 2k\pi}{n} \right) \right]$$

for  $k = 0, 1, \dots, n - 1$ .

These are equally spaced points on the circle of radius  $r^{1/n}$ .

**Example:** The cube roots of  $-8$ : Since  $-8 = 8(\cos(\pi) + i \sin(\pi))$ , we have

$$(-8)^{1/3} = 8^{1/3} \left[ \cos \left( \frac{\pi + 2k\pi}{3} \right) + i \sin \left( \frac{\pi + 2k\pi}{3} \right) \right]$$

for  $k = 0, 1, 2$ . We get

$$2(\cos(\pi/3) + i \sin(\pi/3)) = 2(1/2 + i\sqrt{3}/2) = 1 + \sqrt{3}i$$

$$2(\cos(3\pi/3) + i \sin(3\pi/3)) = 2(-1 + 0i) = -2$$

$$2(\cos(5\pi/3) + i \sin(5\pi/3)) = 2(1/2 - i\sqrt{3}/2) = 1 - \sqrt{3}i$$

## Euler's formula

Using some Calculus, one can prove:

**Theorem (Euler's formula):** For any real number  $x$ ,

$$e^{ix} = \cos x + i \sin x$$

Thus  $e^{ix}$  is a complex number on the unit circle. This is most often used as a shorthand:

$$z = r(\cos \theta + i \sin \theta) = re^{i\theta}$$

It also leads to one of the most remarkable formulas in mathematics, which combines 5 of the most important numbers:

$$e^{i\pi} + 1 = 0$$