

Math 1600 Lecture 33, Section 2, 24 Nov 2014

Announcements:

Today we finish 5.1 and start 5.2. Continue **reading** Section 5.2 for next class, and start reading 5.3. Work through recommended [homework questions](#).

Tutorials: Quiz 9 covers 4.6, 5.1 and the first part of 5.2 (orthogonal complements).

Office hour: Mon 3:00-3:30, MC103B.

Help Centers: Monday-Friday 2:30-6:30 in MC 106.

T/F: A matrix with orthogonal columns is called an orthogonal matrix.

T/F: An orthogonal matrix must be square.

Questions: Why are orthonormal bases great? Are orthogonal bases great too?

Review of Section 5.1: Orthogonal and Orthonormal sets

Definition: A set of vectors $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k\}$ in \mathbb{R}^n is an **orthogonal set** if $\vec{v}_i \cdot \vec{v}_j = 0$ for $i \neq j$. If it is an **orthonormal set** if in addition $\vec{v}_i \cdot \vec{v}_i = 1$ for each i , i.e., each vector is a unit vector.

Theorem 5.1: An orthogonal set of nonzero vectors is always linearly independent.

Definition: An **orthogonal basis** for a subspace W of \mathbb{R}^n is a basis of W that is an orthogonal set. An **orthonormal basis** is a basis that is an orthonormal set.

You only need to check that the set spans W , since it is automatically linearly independent.

Note: An orthogonal basis can be converted to an orthonormal basis by dividing each vector by its length. We'll show in Section 5.3 that every subspace *has* an orthogonal basis.

Recall that if $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k\}$ is any basis of a subspace W , then any \vec{w} in W can be written uniquely as a linearly combination of the vectors in the basis. In general, finding the coefficients involves solving a linear system. For an orthogonal basis, it is much easier:

Theorems 5.2/5.3: If $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k\}$ is an orthogonal basis of a subspace W , and \vec{w} is in W , then

$$\vec{w} = c_1\vec{v}_1 + \dots + c_k\vec{v}_k \quad \text{where} \quad c_i = \frac{\vec{w} \cdot \vec{v}_i}{\vec{v}_i \cdot \vec{v}_i}$$

If the basis is orthonormal, then

$$c_i = \vec{w} \cdot \vec{v}_i$$

Orthogonal Matrices

Definition: A **square** matrix Q **with real entries** whose columns form an orthonormal set is called an **orthogonal** matrix!

Note: In \mathbb{R}^2 and \mathbb{R}^3 , orthogonal matrices correspond exactly to the rotations and reflections. This is an important geometric reason to study them. Another reason is that we will see in Section 5.4 that they are related to diagonalization of symmetric matrices.

Theorems 5.4 and 5.5: Q is orthogonal if and only if $Q^T Q = I$, i.e. if and only if Q is invertible and $Q^{-1} = Q^T$.

Examples: $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$, $B = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ and

$$C = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

Theorem 5.7: If Q is orthogonal, then its **rows** form an orthonormal set too.

Another way to put it is that Q^T is also an orthogonal matrix.

Theorem 5.6: Let Q be an $n \times n$ matrix. Then the following statements are equivalent:

- Q is orthogonal.
- $\|Q\vec{x}\| = \|\vec{x}\|$ for every \vec{x} in \mathbb{R}^n .
- $Q\vec{x} \cdot Q\vec{y} = \vec{x} \cdot \vec{y}$ for every \vec{x} and \vec{y} in \mathbb{R}^n .

New material

Example: Compute the eigenvalues and determinants of C and A on the board.

Theorem 5.8: Let Q be an orthogonal matrix. Then:

- Q^{-1} is orthogonal.
- $\det Q = \pm 1$
- If λ is a real or complex eigenvalue of Q , then $|\lambda| = 1$.
- If Q_1 and Q_2 are orthogonal matrices of the same size, then $Q_1 Q_2$ is orthogonal.

Proof:

(a) is Theorem 5.7, since $Q^{-1} = Q^T$.

(b): Since $I = Q^T Q$, we have

$$1 = \det I = \det(Q^T Q) = \det(Q^T) \det(Q) = \det(Q)^2.$$

Therefore $\det(Q) = \pm 1$.

(c) If $Q\vec{v} = \lambda\vec{v}$, then

$$\|\vec{v}\| = \|Q\vec{v}\| = \|\lambda\vec{v}\| = |\lambda| \|\vec{v}\|$$

so $|\lambda| = 1$, since $\|\vec{v}\| \neq 0$.

(d) Exercise, using properties of transpose. \square

Section 5.2: Orthogonal Complements and Orthogonal Projections

We saw in Section 5.1 that orthogonal and orthonormal bases are particularly easy to work with. In Section 5.3, we will learn how to find these kinds of bases. In this section, we learn the tools which will be needed in Section 5.3. We will also find a new way to understand the subspaces associated to a matrix.

Orthogonal Complements

If W is a plane through the origin, with normal vector \vec{n} , then the subspaces W and $\text{span}(\vec{n})$ have the property that every vector in one is orthogonal to every vector in the other.

Definition: Let W be a subspace of \mathbb{R}^n . A vector \vec{v} is **orthogonal** to W if \vec{v} is orthogonal to every vector in W . The **orthogonal complement** of W is the set of all vectors orthogonal to W and is denoted W^\perp . So

$$W^\perp = \{\vec{v} \in \mathbb{R}^n : \vec{v} \cdot \vec{w} = 0 \text{ for all } \vec{w} \text{ in } W\}$$

In the example above, if we write $\ell = \text{span}(\vec{n})$ for the line perpendicular to W , then $\ell = W^\perp$ and $W = \ell^\perp$.

Theorem 5.9: Let W be a subspace of \mathbb{R}^n . Then:

- W^\perp is a subspace of \mathbb{R}^n .
- $(W^\perp)^\perp = W$
- $W \cap W^\perp = \{\vec{0}\}$
- If $W = \text{span}(\vec{w}_1, \dots, \vec{w}_k)$, then \vec{v} is in W^\perp if and only if $\vec{v} \cdot \vec{w}_i = 0$ for all i .

Explain (a), (c), (d) on board. (b) will be Corollary 5.12.

Theorem 5.10: Let A be an $m \times n$ matrix. Then

$$(\text{row}(A))^\perp = \text{null}(A) \quad \text{and} \quad (\text{col}(A))^\perp = \text{null}(A^T)$$

The first two are in \mathbb{R}^n and the last two are in \mathbb{R}^m . These are the **four**

fundamental subspaces of A .

Let's see why $(\text{row}(A))^\perp = \text{null}(A)$. A vector is in $\text{null}(A)$ exactly when it is orthogonal to the rows of A . But the rows of A span $\text{row}(A)$, so the vectors in $\text{null}(A)$ are exactly those which are orthogonal to $\text{row}(A)$, by 5.9(d).

The fact that $(\text{col}(A))^\perp = \text{null}(A^T)$ follows by replacing A with A^T .

Example: Let W be the subspace spanned by $\vec{v}_1 = [1, 2, 3]$ and $\vec{v}_2 = [2, 5, 7]$. Find a basis for W^\perp .

Solution: Let A be the matrix with \vec{v}_1 and \vec{v}_2 as rows. Then $W = \text{row}(A)$, so $W^\perp = \text{null}(A)$. Continue on board.

Exercise: Let W be the subspace spanned by $\vec{v}_1 = [1, 2, 3]$. Find a basis for W^\perp .

Could use the same method. Or could just find two linearly independent vectors perpendicular to \vec{v}_1 by inspection.

Orthogonal projection

Recall (from waaaaay back in Section 1.2) that the formula for the projection of a vector \vec{v} onto a nonzero vector \vec{u} is:

$$\text{proj}_{\vec{u}}(\vec{v}) = \left(\frac{\vec{u} \cdot \vec{v}}{\vec{u} \cdot \vec{u}} \right) \vec{u}.$$

(Illustrated by this [java applet](#), where red is \vec{v} , blue is \vec{u} and yellow is the projection.)

We didn't name it then, but we also noticed that $\vec{v} - \text{proj}_{\vec{u}}(\vec{v})$ is orthogonal to \vec{u} . Let's call this $\text{perp}_{\vec{u}}(\vec{v})$.

So if we write $W = \text{span}(\vec{u})$, then $\vec{w} = \text{proj}_{\vec{u}}(\vec{v})$ is in W , $\vec{w}^\perp = \text{perp}_{\vec{u}}(\vec{v})$ is in W^\perp , and $\vec{v} = \vec{w} + \vec{w}^\perp$. We can do this more generally:



Definition: Let W be a subspace of \mathbb{R}^n and let $\{\vec{u}_1, \dots, \vec{u}_k\}$ be an **orthogonal** basis for W . For \vec{v} in \mathbb{R}^n , the **orthogonal projection** of \vec{v} onto W is the vector

$$\text{proj}_W(\vec{v}) = \text{proj}_{\vec{u}_1}(\vec{v}) + \dots + \text{proj}_{\vec{u}_k}(\vec{v})$$

The **component of \vec{v} orthogonal to W** is the vector

$$\text{perp}_W(\vec{v}) = \vec{v} - \text{proj}_W(\vec{v})$$

We will show soon that $\text{perp}_W(\vec{v})$ is in W^\perp .

Note that multiplying \vec{u} by a scalar in the earlier example doesn't change W , \vec{w} or \vec{w}^\perp . We'll see later that the general definition also doesn't depend on the choice of orthogonal basis.




Figure 5.8

$$\mathbf{p} = \mathbf{p}_1 + \mathbf{p}_2$$




Figure 5.9

$$\mathbf{v} = \text{proj}_W(\mathbf{v}) + \text{perp}_W(\mathbf{v})$$

Example: Let $W = \text{span}(\vec{u}_1, \vec{u}_2)$, where $\vec{u}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}$ and $\vec{u}_2 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$.

Compute $\text{proj}_W(\vec{v})$ and $\text{perp}_W(\vec{v})$, where $\vec{v} = \begin{bmatrix} 1 \\ 3 \\ 2 \\ 4 \end{bmatrix}$. On board.

Notice that $\text{perp}_W(\vec{v})$ is in W^\perp .