

Math 1600B Lecture 34, Section 2, 31 March 2014

Announcements:

Today we finish 5.2 and start 5.3. **Read** Sections 5.3 and 5.4 for next class. Work through recommended [homework questions](#).

Tutorials: Review and questions.

Office hour: Mon 1:30-2:30 and Wed 10:30-11:15, MC103B.

Help Centers: Monday-Friday 2:30-6:30 in MC 106.

Final exam: Covers whole course, with an emphasis on the material after the midterm. Our course will end with Section 5.4.

Question: If $W = \mathbb{R}^n$, then $W^\perp = \{\vec{0}\}$

T/F: An orthogonal basis $\{\vec{v}_1, \dots, \vec{v}_k\}$ must have

$$\vec{v}_i \cdot \vec{v}_j = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases}$$

Review of Section 5.2: Orthogonal Complements and Orthogonal Projections

We saw in Section 5.1 that orthogonal and orthonormal bases are particularly easy to work with. In Section 5.3 (today), we will learn how to find these kinds of bases. In Section 5.2, we learn the tools which will be needed in Section 5.3.

Orthogonal Complements

Definition: Let W be a subspace of \mathbb{R}^n . A vector \vec{v} is **orthogonal** to W if \vec{v} is orthogonal to every vector in W . The **orthogonal complement** of W is the set of all vectors orthogonal to W and is denoted W^\perp . So

$$W^\perp = \{\vec{v} \in \mathbb{R}^n : \vec{v} \cdot \vec{w} = 0 \text{ for all } \vec{w} \text{ in } W\}$$

An example to keep in mind is where W is a plane through the origin in \mathbb{R}^3 and W^\perp is $\text{span}(\vec{n})$, where \vec{n} is the normal vector to W .

Theorem 5.9: Let W be a subspace of \mathbb{R}^n . Then:

- W^\perp is a subspace of \mathbb{R}^n .
- $(W^\perp)^\perp = W$
- $W \cap W^\perp = \{\vec{0}\}$
- If $W = \text{span}(\vec{w}_1, \dots, \vec{w}_k)$, then \vec{v} is in W^\perp if and only if $\vec{v} \cdot \vec{w}_i = 0$ for all i .

We proved all of these except part (b), which will come today.

Theorem 5.10: Let A be an $m \times n$ matrix. Then

$$(\text{row}(A))^\perp = \text{null}(A) \quad \text{and} \quad (\text{col}(A))^\perp = \text{null}(A^T)$$

The first two are in \mathbb{R}^n and the last two are in \mathbb{R}^m . These are the **four fundamental subspaces** of A .

Orthogonal projection


Let \vec{u} be a nonzero vector in \mathbb{R}^n , and for any \vec{v} in \mathbb{R}^n define:

$$\text{proj}_{\vec{u}}(\vec{v}) = \left(\frac{\vec{u} \cdot \vec{v}}{\vec{u} \cdot \vec{u}} \right) \vec{u}.$$

$$\text{perp}_{\vec{u}}(\vec{v}) = \vec{v} - \text{proj}_{\vec{u}}(\vec{v})$$

If we write $W = \text{span}(\vec{u})$, then $\vec{w} = \text{proj}_{\vec{u}}(\vec{v})$ is in W , $\vec{w}^\perp = \text{perp}_{\vec{u}}(\vec{v})$ is in W^\perp , and $\vec{v} = \vec{w} + \vec{w}^\perp$. We can do this more generally:

Definition: Let W be a subspace of \mathbb{R}^n and let $\{\vec{u}_1, \dots, \vec{u}_k\}$ be an orthogonal basis for W . For \vec{v} in \mathbb{R}^n , the **orthogonal projection** of \vec{v} onto W is the vector



$$\text{proj}_W(\vec{v}) = \text{proj}_{\vec{u}_1}(\vec{v}) + \cdots + \text{proj}_{\vec{u}_k}(\vec{v})$$

The **component of \vec{v} orthogonal to W** is the vector

$$\text{perp}_W(\vec{v}) = \vec{v} - \text{proj}_W(\vec{v})$$

We will show soon that $\text{perp}_W(\vec{v})$ is in W^\perp .

Note that multiplying \vec{u} by a scalar in the earlier example doesn't change W , \vec{w} or \vec{w}^\perp . We'll see later that the general definition also doesn't depend on the choice of orthogonal basis.

New material

Theorem: $\text{perp}_W(\vec{v})$ is in W^\perp .

Explain on board.

Now we will see that proj and perp don't depend on the choice of orthogonal basis. Here and in the rest of the section, we **assume that every subspace has at least one orthogonal basis**.

Theorem 5.11: Let W be a subspace of \mathbb{R}^n and let \vec{v} be a vector in \mathbb{R}^n . Then there are **unique** vectors \vec{w} in W and \vec{w}^\perp in W^\perp such that $\vec{v} = \vec{w} + \vec{w}^\perp$.

Proof: We saw above that such a decomposition exists, by taking $\vec{w} = \text{proj}_W(\vec{v})$ and $\vec{w}^\perp = \text{perp}_W(\vec{v})$, using an orthogonal basis for W .

We now show that this decomposition is unique. So suppose $\vec{v} = \vec{w}_1 + \vec{w}_1^\perp$ is another such decomposition. Then $\vec{w} + \vec{w}^\perp = \vec{w}_1 + \vec{w}_1^\perp$, so

$$\vec{w} - \vec{w}_1 = \vec{w}_1^\perp - \vec{w}^\perp$$

The left hand side is in W and the right hand side is in W^\perp (why?), so both sides must be zero (why?). So $\vec{w} = \vec{w}_1$ and $\vec{w}^\perp = \vec{w}_1^\perp$. \square



Figure 5.8

$$\mathbf{p} = \mathbf{p}_1 + \mathbf{p}_2$$




Figure 5.9

$$\mathbf{v} = \text{proj}_W(\mathbf{v}) + \text{perp}_W(\mathbf{v})$$

Note that \perp is an operation on subspaces, but is not an operation on vectors.

Now we can prove part (b) of Theorem 5.9.

Corollary 5.12: If W is a subspace of \mathbb{R}^n , then $(W^\perp)^\perp = W$.

Proof: If \vec{w} is in W and \vec{x} is in W^\perp , then $\vec{w} \cdot \vec{x} = 0$. This means that \vec{w} is in $(W^\perp)^\perp$. So $W \subseteq (W^\perp)^\perp$.

We need to show that every vector in $(W^\perp)^\perp$ is in W . So let \vec{v} be a vector in $(W^\perp)^\perp$. By the previous result, we can write \vec{v} as $\vec{w} + \vec{w}^\perp$, where \vec{w} is in W and \vec{w}^\perp is in W^\perp . Then

$$\begin{aligned} 0 &= \vec{v} \cdot \vec{w}^\perp = (\vec{w} + \vec{w}^\perp) \cdot \vec{w}^\perp \\ &= \vec{w} \cdot \vec{w}^\perp + \vec{w}^\perp \cdot \vec{w}^\perp = 0 + \vec{w}^\perp \cdot \vec{w}^\perp = \vec{w}^\perp \cdot \vec{w}^\perp \end{aligned}$$

So $\vec{w}^\perp = \vec{0}$ and $\vec{v} = \vec{w}$ is in W . \square

This next result is related to the Rank Theorem:

Theorem 5.13: If W is a subspace of \mathbb{R}^n , then

$$\dim W + \dim W^\perp = n$$

Proof: Let $\{\vec{u}_1, \dots, \vec{u}_k\}$ be an orthogonal basis of W and let $\{\vec{v}_1, \dots, \vec{v}_\ell\}$ be an orthogonal basis of W^\perp . Then $\{\vec{u}_1, \dots, \vec{u}_k, \vec{v}_1, \dots, \vec{v}_\ell\}$ is an orthogonal basis for \mathbb{R}^n . (Explain.) The result follows. \square

Example: For W a plane in \mathbb{R}^3 , $2 + 1 = 3$.

The Rank Theorem follows if we take $W = \text{row}(A)$, since then $W^\perp = \text{null}(A)$:

Corollary 5.14 (The Rank Theorem, again): If A is an $m \times n$ matrix, then

$$\text{rank}(A) + \text{nullity}(A) = n$$

Note: The logic here can be reversed. We can use the rank theorem to prove Theorem 5.13, and Theorem 5.13 can be used to prove Corollary 5.12.

Section 5.3: The Gram-Schmidt Process and the QR Factorization

The Gram-Schmidt Process

This is a fancy name for a way of converting a basis into an orthogonal or orthonormal basis. And it's pretty clear how to do it, given what we know.

Example: Let $W = \text{span}(\vec{x}_1, \vec{x}_2)$ where $\vec{x}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ and $\vec{x}_2 = \begin{bmatrix} -2 \\ 0 \\ 1 \end{bmatrix}$.

Find an orthogonal basis for W .

Solution: Ideas? Do on board.

Question: What if we had a third basis vector \vec{x}_3 ?

Theorem 5.15 (The Gram-Schmidt Process): Let $\{\vec{x}_1, \dots, \vec{x}_k\}$ be a basis for a subspace W of \mathbb{R}^n . Write $W_1 = \text{span}(\vec{x}_1)$, $W_2 = \text{span}(\vec{x}_1, \vec{x}_2)$, \dots , $W_k = \text{span}(\vec{x}_1, \dots, \vec{x}_k)$. Define:

$$\vec{v}_1 = \vec{x}_1$$

$$\vec{v}_2 = \text{perp}_{W_1}(\vec{x}_2) = \vec{x}_2 - \frac{\vec{v}_1 \cdot \vec{x}_2}{\vec{v}_1 \cdot \vec{v}_1} \vec{v}_1$$

$$\vec{v}_3 = \text{perp}_{W_2}(\vec{x}_3) = \vec{x}_3 - \frac{\vec{v}_1 \cdot \vec{x}_3}{\vec{v}_1 \cdot \vec{v}_1} \vec{v}_1 - \frac{\vec{v}_2 \cdot \vec{x}_3}{\vec{v}_2 \cdot \vec{v}_2} \vec{v}_2$$

\vdots

$$\vec{v}_k = \text{perp}_{W_{k-1}}(\vec{x}_k) = \vec{x}_k - \frac{\vec{v}_1 \cdot \vec{x}_k}{\vec{v}_1 \cdot \vec{v}_1} \vec{v}_1 - \dots - \frac{\vec{v}_{k-1} \cdot \vec{x}_k}{\vec{v}_{k-1} \cdot \vec{v}_{k-1}} \vec{v}_{k-1}$$

Then for each i , $\{\vec{v}_1, \dots, \vec{v}_i\}$ is an orthogonal basis for W_i . In particular, $\{\vec{v}_1, \dots, \vec{v}_k\}$ is an orthogonal basis for $W = W_k$.

Explain verbally.

Notes: To compute perp_{W_i} you have to use the *orthogonal* basis of \vec{v}_j 's that you have constructed already, not the original basis of \vec{x}_j 's.

The basis you get depends on the order of the vectors you start with. You should always do a question using the vectors in the order given, since that order will be chosen to minimize the arithmetic.

If you are asked to find an orthonormal basis, normalize each \vec{v}_j at the end. (It is correct to normalize earlier, but can be messier.)